



Heat effects on mortality in 15 European cities

Author(s): Baccini M, Biggeri A, Accetta G, Kosatsky T, Katsouyanni K, Analitis A, anderson HR, Bisanti L, D'Ippoliti D, Danova J, Forsberg B, Medina S, Paldy A, Rabaczko D, Schindler C, Michelozzi P

Year: 2008

Journal: Epidemiology. 19 (5): 711-719

Abstract:

Background: Epidemiologic studies show that high temperatures are related to mortality, but little is known about the exposure-response function and the lagged effect of heat. We report the associations between daily maximum apparent temperature and daily deaths during the warm season in 15 European cities. Methods: The city-specific analyses were based on generalized estimating equations and the city-specific results were combined in a Bayesian random effects meta-analysis. We specified distributed lag models in studying the delayed effect of exposure. Time-varying coefficient models were used to check the assumption of a constant heat effect over the warm season. Results: The city-specific exposure-response functions have a V shape, with a change-point that varied among cities. The meta-analytic estimate of the threshold was 29.4°C for Mediterranean cities and 23.3°C for north-continental cities. The estimated overall change in all natural mortality associated with a 1°C increase in maximum apparent temperature above the city-specific threshold was 3.12% (95% credibility interval Euro Surveillance (Bulletin European Sur Les Maladies Transmissibles; European Communicable Disease Bulletin) 0.60% to 5.72%) in the Mediterranean region and 1.84% (0.06% to 3.64%) in the north-continental region. Stronger associations were found between heat and mortality from respiratory diseases, and with mortality in the elderly. Conclusions: There is an important mortality effect of heat across Europe. The effect is evident from June through August; it is limited to the first week following temperature excess, with evidence of mortality displacement. There is some suggestion of a higher effect of early season exposures. Acclimatization and individual susceptibility need further investigation as possible explanations for the observed heterogeneity among cities.

Source:

http://journals.lww.com/epidem/Fulltext/2008/09000/Heat_Effects_on_Mortality_in_15_European_Cities.13.aspx

Resource Description

Exposure :

weather or climate related pathway by which climate change affects health

Temperature

Temperature: Extreme Heat

Geographic Feature:

resource focuses on specific type of geography

None or Unspecified

Climate Change and Human Health Literature Portal

Geographic Location: ☒

resource focuses on specific location

Non-United States

Non-United States: Europe

European Region/Country: European Country

Other European Country : Greece;Spain;Hungary;Ireland;Finland;Slovenia;England;Italy;France;Czech Republic;Sweden;Switzerland

Health Impact: ☒

specification of health effect or disease related to climate change exposure

Injury

Mitigation/Adaptation: ☒

mitigation or adaptation strategy is a focus of resource

Adaptation

Population of Concern: A focus of content

Population of Concern: ☒

populations at particular risk or vulnerability to climate change impacts

Elderly, Low Socioeconomic Status

Resource Type: ☒

format or standard characteristic of resource

Research Article

Timescale: ☒

time period studied

Time Scale Unspecified

Vulnerability/Impact Assessment: ☒

resource focus on process of identifying, quantifying, and prioritizing vulnerabilities in a system

A focus of content